

## INDEX TO VOLUME XXIII

New names and the final members of new combinations are in **bold face type**. Mistakes in the spelling of scientific names encountered in text are corrected in the index.

- Abutilon sylvaticum*, 480  
*Acacia*, 335–337; *angustissima*, 337;  
     *Farnesiana*, 335; *pedicellata*, 338;  
     *riparia*, 335  
*Acaulium*, 314–317, 323, 325, 326,  
     329; *albo-nigrescens*, 316, 317, 324,  
     326, 329  
*Acer*, 118, 120, 131, 303  
*Achlya*, 58, 309  
*Achorion gypseum*, 87, 88, 90, 92–94  
*Achyranthes bidentata*, 378; *biden-*  
     *tata japonica*, 378; *crispa*, 378  
*Actinomyces scabies*, 304  
*Actinostemon*, 467; *concolor*, 470  
*Adesmia elegans*, 351; *laxa*, 351; *punc-*  
     *tata*, 348  
*Aecidiolum Hydrocotyles*, 486  
*Aecidium*, 101, 235, 360, 463, 473,  
     475, 477, 486, 496; *Azorellae*, 487;  
     *brasiliense*, 500; *ceraceum*, 492; *circ-*  
     *umscriptum*, 475; *Cissi*, 475; *Cordi-*  
     *ae*, 500; *desmum*, 478; *Desmodii*,  
     344; *distinguendum*, 497; *Gayo-*  
     *phyti*, 485; *Glaziovii*, 483; *gutta-*  
     *tum*, 475; *Hieronymi*, 335; *hydro-*  
     *cotylum*, 486; *Hyperici-frondosi*,  
     481; *hypsophilum*, 463, 470; *Ipo-*  
     *moeae*, 498; *lathyrinum*, 354; *Lind-*  
     *davianum*, 500; *Malvastri*, 478;  
     *Maublancii*, 359; *mexicanum*, 359;  
     *Nectandrae*, 102; *ochraceum*, 492;  
     *Oenotherae*, 485; *Opuntiae*, 482;  
     *Pereskiae*, 472, 473; *Philippii*, 489;  
     *punctatum*, 104; *Rickii*, 364; *Sphae-*  
     *ralceae*, 478; *Tournefortiae*, 500;  
     *Trifolii-megalanthi*, 356; *Triumfet-*  
     *tae*, 477; ***vinnulum***, 360; *Violae*,  
     482; *xanthoxylum*, 364  
*Agaricus*, 240  
*Ageratum conyzoides*, 402  
*Agoseris*, 80; *aurantiaca*, 80; *humilis*,  
     80; *parviflora*, 80; *scorzonerifolia*,  
     80  
*Agrimonia hirsuta*, 104; *striata*, 79  
*Albugo*, 201, 255; *candida*, 304  
*Alchornea Iricurana*, 466; *pyncogyne*,  
     466  
*Allium brevistylum*, 82; *Cepa*, 303  
*Allodus interveniens*, 478; *opulenta*,  
     498  
***Allopuccinia***, 347; ***diluta***, 347, 348, 349  
*Alnus*, 118, 120; *glutinosa*, 147; *in-*  
     *cana*, 301  
*Alternaria*, 160, 178–185; *Brassicae*,  
     179, 180, 302; *Solani*, 180, 183–185,  
     304; *Solani* ***Symphoricarpi***, 178,  
     184, 185; *tenuis*, 178, 183, 184  
*Althaea*, 479, 480; *rosea*, 81  
*Alveolaria Cordiae*, 501  
*Amanita*, 225; *caesarea*, 225; *calyp-*  
     *trata*, 225, 226; *calyptroides*, 225;  
     *calyptroderma*, 225, 226  
*Amanita calypttrata* and *Amanita*  
     *calyptroderma*, S. M. Zeller, 225  
*Amaranthus*, 390  
*Ambrosia trifida*, 393  
*Amicia*, 348; *Lobbiana*, 348; *parvula*,  
     357  
*Amorpha canescens*, 81  
*Amorphophallus*, 219  
*Amygdalus Persica*, 303  
*Ananas sativus*, 272, 273, 275, 277,  
     279–281, 283, 284, 286, 290  
*Andropogon leucostachyus*, 297  
*Angelica Grayi*, 80  
*Anthracnose*, *alternariose* and *Botry-*  
     *tis* rot of the snowberry, W. H.  
     Davis, 159  
*Anticlea coloradensis*, 79  
*Antirrhinum majus*, 304  
*Aphragmium*, 201, 253, 254  
*Aplanes*, 309  
*Aplanobacter insidiosus*, 301  
*Arachis hypogaea*, 303, 369  
*Araiospora*, 201  
*Arauja sericifera*, 493  
*Arctium Lappa*, 393  
*Arracacia xanthorrhiza*, 490  
*Artemisia cana*, 79; *cuneata*, 79  
***Arthuria***, 463; ***catenulata***, 464  
*Ascigerous* stage of *Colletotrichum*  
     *lagenarium* induced by ultra-violet  
     irradiation, F. L. Stevens, 134  
*Asclepias*, 494; *curassavica*, 493, 494  
*Ascobelus*, 248; *magnificus*, 33, 35, 47;  
     *Winteri*, 34  
*Ascocarpic* stage of species of *Scopu-*  
     *lariopsis*, C. W. Emmons and B. O.  
     Dodge, 313  
*Ascochyta*, 302; *Gossypii*, 302

## INDEX TO VOLUME XXIII

New names and the final members of new combinations are in **bold face type**. Mistakes in the spelling of scientific names encountered in text are corrected in the index.

- Abutilon sylvaticum*, 480  
*Acacia*, 335–337; *angustissima*, 337;  
     *Farnesiana*, 335; *pedicellata*, 338;  
     *riparia*, 335  
*Acaulium*, 314–317, 323, 325, 326,  
     329; *albo-nigrescens*, 316, 317, 324,  
     326, 329  
*Acer*, 118, 120, 131, 303  
*Achlya*, 58, 309  
*Achorion gypseum*, 87, 88, 90, 92–94  
*Achyranthes bidentata*, 378; *biden-*  
     *tata japonica*, 378; *crispa*, 378  
*Actinomyces scabies*, 304  
*Actinostemon*, 467; *concolor*, 470  
*Adesmia elegans*, 351; *laxa*, 351; *punc-*  
     *tata*, 348  
*Aecidiolum Hydrocotyles*, 486  
*Aecidium*, 101, 235, 360, 463, 473,  
     475, 477, 486, 496; *Azorellae*, 487;  
     *brasiliense*, 500; *ceraceum*, 492; *circ-*  
     *umscriptum*, 475; *Cissi*, 475; *Cordi-*  
     *ae*, 500; *desmum*, 478; *Desmodii*,  
     344; *distinguendum*, 497; *Gayo-*  
     *phyti*, 485; *Glaziovii*, 483; *gutta-*  
     *tum*, 475; *Hieronymi*, 335; *hydro-*  
     *cotylum*, 486; *Hyperici-frondosi*,  
     481; *hypsophilum*, 463, 470; *Ipo-*  
     *moeae*, 498; *lathyrinum*, 354; *Lind-*  
     *davianum*, 500; *Malvastri*, 478;  
     *Maublancii*, 359; *mexicanum*, 359;  
     *Nectandrae*, 102; *ochraceum*, 492;  
     *Oenotherae*, 485; *Opuntiae*, 482;  
     *Pereskiae*, 472, 473; *Philippii*, 489;  
     *punctatum*, 104; *Rickii*, 364; *Sphae-*  
     *ralceae*, 478; *Tournefortiae*, 500;  
     *Trifolii-megalanthi*, 356; *Triumfet-*  
     *tae*, 477; ***vinnulum***, 360; *Violae*,  
     482; *xanthoxylum*, 364  
*Agaricus*, 240  
*Ageratum conyzoides*, 402  
*Agoseris*, 80; *aurantiaca*, 80; *humilis*,  
     80; *parviflora*, 80; *scorzonerifolia*,  
     80  
*Agrimonia hirsuta*, 104; *striata*, 79  
*Albugo*, 201, 255; *candida*, 304  
*Alchornea Iricurana*, 466; *pyncogyne*,  
     466  
*Allium brevistylum*, 82; *Cepa*, 303  
*Allodus interveniens*, 478; *opulenta*,  
     498  
***Allopuccinia***, 347; ***diluta***, 347, 348, 349  
*Alnus*, 118, 120; *glutinosa*, 147; *in-*  
     *cana*, 301  
*Alternaria*, 160, 178–185; *Brassicae*,  
     179, 180, 302; *Solani*, 180, 183–185,  
     304; *Solani* ***Symphoricarpi***, 178,  
     184, 185; *tenuis*, 178, 183, 184  
*Althaea*, 479, 480; *rosea*, 81  
*Alveolaria Cordiae*, 501  
*Amanita*, 225; *caesarea*, 225; *calyp-*  
     *trata*, 225, 226; *calyptroides*, 225;  
     *calyptroderma*, 225, 226  
*Amanita calyptrata* and *Amanita*  
     *calyptroderma*, S. M. Zeller, 225  
*Amaranthus*, 390  
*Ambrosia trifida*, 393  
*Amicia*, 348; *Lobbiana*, 348; *parvula*,  
     357  
*Amorpha canescens*, 81  
*Amorphophallus*, 219  
*Amygdalus Persica*, 303  
*Ananas sativus*, 272, 273, 275, 277,  
     279–281, 283, 284, 286, 290  
*Andropogon leucostachyus*, 297  
*Angelica Grayi*, 80  
*Anthracnose*, *alternariose* and *Botry-*  
     *tis* rot of the snowberry, W. H.  
     Davis, 159  
*Anticlea coloradensis*, 79  
*Antirrhinum majus*, 304  
*Aphragmium*, 201, 253, 254  
*Aplanes*, 309  
*Aplanobacter insidiosus*, 301  
*Arachis hypogaea*, 303, 369  
*Araiospora*, 201  
*Arauja sericifera*, 493  
*Arctium Lappa*, 393  
*Arracacia xanthorrhiza*, 490  
*Artemisia cana*, 79; *cuneata*, 79  
***Arthuria***, 463; ***catenulata***, 464  
*Ascigerous* stage of *Colletotrichum*  
     *lagenarium* induced by ultra-violet  
     irradiation, F. L. Stevens, 134  
*Asclepias*, 494; *curassavica*, 493, 494  
*Ascobelus*, 248; *magnificus*, 33, 35, 47;  
     *Winteri*, 34  
*Ascocarpic* stage of species of *Scopu-*  
     *lariopsis*, C. W. Emmons and B. O.  
     Dodge, 313  
*Ascochyta*, 302; *Gossypii*, 302

- Ascoidea, 51-55, 58, 61, 67-69, 75;  
rubescens, 51, 59, 65, 68, 69, 75;  
saprolegnoides, 52, 68  
Asclepias disciflora, 104  
Ascotricha, 323  
Asimina, 120  
Aspergillus, 75, 89, 313, 323, 326, 329;  
herbariorum, 326  
Aster, 79  
Astronium, 472  
Avena sativa, 265, 303  
  
Bacillus amylovorus, 301, 303; carotovor-  
us, 303, 304; phytophthorus,  
304; prodigiosus, 140; tracheiphilus,  
302  
Bacterium campestre, 304; Citri, 302;  
Erodii, 303; Phaseoli, 302; Pruni,  
303; Sojae, 304; solanacearum, 303;  
tumefaciens, 303, 304  
Bailey, F. D., and Zeller, S. M., The  
occurrence of Schizophyllum com-  
mune on green apples, 154  
Banisteria campestris, 362  
Bartholomew, E. T., Herbarium ar-  
rangement of mycological speci-  
mens, 227  
Basidiophora, 255  
Bauhinia, 343, 344; forficata, 344;  
forficata latifolia, 344; rufa, 344  
Berberis, 98, 99, 101; Darwinii, 96;  
divaricata, 100; glaucescens, 99;  
phyllacantha, 99  
Beta vulgaris, 286  
Betula, 120  
Bignonia capreolata, 302  
Bilbergia, 281  
Bistorta vivipara, 81  
Bixa orellana, 376  
Blain, W. L., A list of diseases of eco-  
nomic plants in Alabama, 300  
Blumenbachia, 482; lateritia, 482  
Boisduvalia concinna, 485  
Bommerella, 322; trigonospora, 322,  
323  
Borreria, 385; micrantha, 385  
Botrytis, 160, 186, 187, 313; Allii,  
303; furcata, 188; Paeoniae, 303;  
vulgaris, 186-188  
Bouteloua simplex, 482  
Bradburya pubescens, 352, 373, 376;  
virginiana, 352  
Bradyspora, 260, 261, 262, 270, 292  
Brassica, 392; campestris, 392; jun-  
cea, 392; Napus, 392; nigra, 392;  
oleracea, 302; oleracea acephala,  
302; oleracea capitata, 179, 392;  
Rapa, 304, 392  
Bromus, 102; ciliatus, 79  
Bubakia, 464, 469; argentinensis, 465,  
468; Crotonis, 465; mexicana, 470;  
Ulei, 466  
Bullaria Gouaniae, 474; inflata, 361;  
invaginata, 474  
Byrsonima crassifolia, 361; interme-  
dia, 360, 361  
  
Caeoma Agrimoniae, 104; Berberidis,  
96; Eugeniarum, 483; Hydroco-  
tyles, 486  
Cajanus indicus, 273, 276, 277  
Caladium, 397  
Calliandra, 334; falcata, 337; laxa,  
334  
Callistephus chinensis, 301  
Calopogonium, 379; orthocarpum, 401  
Caltha rotundifolia, 81  
Calyptospora columnaris, 78  
Camellia japonica, 303  
Cannabis sativa, 265, 272  
Capsicum annum, 303  
Cardamine cordifolia, 79  
Cardiospermum Halicacabum, 473  
Carex, 78; nebraskensis, 81; Pseudo-  
cyperus, 297; siccata, 80  
Carica Papaya, 266, 270-273, 276,  
283, 284, 286, 290-293  
Carpinus, 120; caroliniana, 302  
Cassia, 332, 341, 342, 343; bicapsu-  
laris, 343; corymbosa, 369; excelsa,  
343; occidentalis, 369; versicolor,  
339  
Castanea, 118  
Castilleja sulphurea, 78  
Cayaponia, 386  
Cejp, K., Notes on Iowa species of  
the genus Irpex, 130  
Cephalanthus, 120  
Ceratostomella fimbriata, 304  
Cercospora, 365, 366, 377, 381, 382,  
402; Achyranthis, 378; Agerati,  
402; alabamensis, 398; althaeina,  
389; althaeina Modiolae, 388; Ama-  
ryllidis, 380; Arachidis, 369; Arcti-  
ambrosiae, 393; Arctii, 393; Atkin-  
sonii, 388; atricincta, 389; biformis,  
380; Bixae, 375; Bloxami, 392; Bol-  
leana, 303; boringuensis, 400; Bor-  
reriae, 385; brachypoda, 395; Brad-  
buryae, 373, 376; Caladii, 397,  
Callae, 397; Calopogonii, 377, 379;  
canescens, 389, 390; capreolata,  
302; Cassavae, 371, 381; Cayapo-  
niae, 365, 386, 401; Cearae, 371-  
382; citrullina, 395; cornicola, 302,  
Costi, 374; cruenta, 302, 304, 384;  
385, 390; cucurbiticola, 386; cylin-  
drospora, 374, 376; depazeoides,  
370, 371; Dolichi, 385; Erechthitis,  
391; flagellaris, 400; fuliginosa, 303;  
fusca, 303; Gilbertii, 404; gonato-  
clada, 403; guianensis, 374, 375;  
Henningsii, 371, 382, 404; Hibisci,  
378, 394; Hibisci-manihotis, 395;

- hibiscina, 379, 382, 395; Hurae, 372; Hymenocallidis, 380; Ipomoeae, 399; **Leonuri**, 395, 396; Ligustri, 304; Lilacis, 303; **malayensis**, 388, 394; Manihotis, 371, 382; Medicaginis, 302; mikaniacola, 397; Modiolae, 389; Nicotianae, 368, 386; pachyspora, 397; Pancratii, 379, 380; penicillata, 370; Perseae, 372; personata, 303, 368, 369; Pipituri, 383; Plantaginis, 391; Porophylli, 365, 393; purpurea, 372; Raciborskii, 368; rigospora, 383; Sambuci, 370, 371; sambucina, 370; Tectoniae, 399; ticinensis, 370; **trinidadensis**, 376, 377; verruculosa, 396, 397
- Cercospora studies—II. Some tropical Cercosporae, W. G. Solheim and F. L. Stevens, 365
- Cerotelium desmium, 478; malvicolium, 478
- Chaetomium, 314, 322, 323, 329; chartarum, 323; pusillum, 323; quadrangulatum, 323; trigonosporum, 323
- Chionothrix, 476
- Chloroscypha**, 248; **chloromela**, 248, 250; **Jacksoni**, 248, 249; **juniperina**, 248, 250; **Seaveri**, 248–250
- Chlorosplenium, 247; chloromelum, 249, 250
- Chrysanthemum, 302
- Chrysocelis Lupini, 345
- Chrysocyclus, 104
- Chrysomyxa, 463
- Ciferri, R., Contributions to the classification of Torulopsidaceae—I. An American variety of the Torulopsis minuta, 140
- Cinnamomum Camphora, 302
- Cintractia, 299; **dubiosa**, 299
- Cirsium, 79; megacephalum, 79
- Cissus sicyoides, 475
- Citrullus vulgaris, 304, 395
- Citrus, 302
- Cladosporium, 402; carpophilum, 303; Citri, 302; effusum, 303; Paeoniae, 303; personatum, 369; Vignae, 302
- Clematis, 302
- Cleome gigantea, 103
- Clitoria, 355; brachystegia, 355; canjanifolia, 355
- Clusia multiflora, 481
- Cocos nucifera, 266
- Coenomyces, 308
- Coleosporium delicatulum, 303, 304; guaranticum, 495; inconspicuum, 304; Ipomoeae, 303, 304, 495; pallidulum, 477; ribicola, 78
- Colletotrichum Carica, 303; gloeosporioides, 302; graminicola, 304; gri-seum, 303; Higginsianum, 304; lagenarium, 134, 135, 137–139, 302; Lindemuthianum, 302; oligochaetum, 139; Pisi, 304; Trifolii, 302
- Colus, 83, 84; javanicus, 83, 84; Schellenbergiae, 83, 84
- Commelina nudiflora, 273
- Comparative study of Sclerotium Rolfsii and Sclerotium Delphinii, F. L. Stevens, 204
- Concept of Mycorrhiza, Arthur Pier-son Kelley, 147
- Condylocarpon Rauwolfiae, 493
- Conidiobolus utriculosus, 309
- Coniothyrium terricola, 314
- Conium maculatum, 488
- Contributions to the classification of Torulopsidaceae—I. An American variety of the Torulopsis minuta, R. Ciferri, 140
- Convolvulus, 495
- Cordia, 500, 501; curassavica, 500; cylindrostachya, 501; discolor, 500; obscura, 500; urticaefolia, 500
- Coriolus, 130
- Cornus, 302
- Corticium Stevensii, 301; vagum, 302, 304
- Costus, 374
- Crataegus succulenta, 78
- Crepis, 80; runcinata, 80
- Crinum americanum, 380
- Cronartium cerebrum, 303; Harknessii, 78; notatum, 361
- Crossopora, 360; notata, 360, 361
- Crotalaria anagyroides, 346; vitellina, 346
- Croton, 377, 464, 469, 470; celtidifolius, 464; chaetophorus, 465; compressus, 468; gossypifolius, 377; hirtus, 465
- Cryptococcus glutinis, 140, 141, 145
- Cucumis Citrullus, 266; Melo, 266, 302; sativus, 200, 290, 302
- Cucurbita maxima, 304; Pepo, 266
- Cummins, George B., Phragmidium species of North America: differential teliospore and aecial characters, 433
- Cuphea dipetala, 483
- Cyathula achyranthoides, 403
- Cylindrosporium Chrysanthemi, 302; Clematidis, 302
- Cystingophora Hieronymi, 335
- Cytospora chrysosperma, 245; nivea, 245
- Dactylomyces, 316; thermophilus, 316
- Daedalea, 130
- Dasiphora fruticosa, 436
- Dasystephana, 79
- Daucus Carota, 267

- Davis, W. H., Anthracnose, alternari-  
ose and Botrytis rot of the snow-  
berry, 159
- Dearness, John, *Volvaria speciosa*,  
152
- Delphinium, 204, 303
- Dendroecia, 338
- Dermatea juniperina, 248, 250
- Derris elliptica, 342
- Desmodium, 344, 352, 355; adscen-  
dens, 355; incanum, 352; uncina-  
tum, 344, 355
- Diagnoses of American Porias—III.  
Some additional brown species,  
with a key to the common brown  
species of the United States and  
Canada, L. O. Overholts, 117
- Dianthus barbatus, 304
- Diaporthe, 223; umbrina, 223, 304
- Dibotryon morbosum, 304
- Dicaeoma montanense, 101
- Dichaena binata, 333; **superba**, 333,  
334
- Dichondra, 496
- Didymaria **borinquensis**, 400; **con-  
jugans**, 401
- Didymopora, 476; **Triumfettae**, 476
- Diorchidium, 333, 348; Berberidis, 99;  
brasilense, 332; insuetum, 362;  
Piptadeniae, 332
- Diospyros Kaki, 303
- Diplocarpon, 447, 451-454, 456-459;  
Rosae, 304, 446
- Dipodascus albidus, 52
- Dispira, 307; americana, 307
- Dodge, B. O., A further study of the  
morphology and life history of the  
rose black spot fungus, 446; In-  
heritance of the albinistic non-  
conidial character in interspecific  
hybrids in Neurospora, 1
- Dodge, B. O., and Emmons, C. W.,  
The ascocarpic stage of species of  
Scopulariopsis, 313
- Dolichos, 384; Lablab, 390
- Earlea, 433-436, 438, 444
- Edythea**, 97; **Berberidis**, 98, 99, 101;  
**quintensis**, 98, 99, 100; **tenella**, 98,  
100, 101, 102
- Elaeagnus, 149
- Elvela, 409; caroliniana, 409; escu-  
lenta, 409; Gigas, 409
- Elymus, 102
- Emmons, C. W., Observations on  
Achorion gypseum, 87
- Emmons, C. W., and Dodge, B. O.,  
The ascocarpic stage of species of  
Scopulariopsis, 313
- Empusa, 414; Muscae, 414; radicans,  
411, 412; sepulchralis, 417
- Endomyces decipiens, 59; fibuliger, 59
- Endophylloides, 476
- Endophyllum, 463, 470, 475, 477, 496;  
circumscription, 475; guttatum, 475
- Entomophthora, 412, 417; americana,  
423; gloeospora, 417, 418; Phyto-  
nomi, 412; Pseudococci, 425; radi-  
cans, 411, 412; sphaerosperma, 411-  
414, 418, 419, 421, 423-428
- Epidermophyton, 314; inguinale, 314
- Epilobium anagallidifolium, 81; an-  
dicolum, 485
- Eragrostis aspera, 299
- Erechtites, 391; praealta, 391
- Eriosema, 355; crinitum, 351
- Erysiphe Polygoni, 302
- Erythrina, 345; Crista-galli, 345
- Erythronium parviflorum, 82
- Erythroxyton Coca, 360
- Eucercospora, 387
- Eugenia, 484
- Euonymus, 303
- Eupatorium repandum, 402
- Euphorbia, 467, 471; heterophylla,  
471; hirta, 471; hypericifolia, 471;  
lasiocarpa, 471; portulacoides, 463,  
470; thymifolia, 471
- Euphragmidium, 433, 434, 436, 438,  
440, 442-444
- Evolvulus sericeus, 497
- Exoascus mirabilis, 304; Pruni, 304
- Exobasidium Vaccinii, 302
- Exosporium concentricum, 303; de-  
pazeoides, 370
- Farysia olivacea, 297; **Pseudocyperi**,  
297
- Ficus Carica, 303
- Fomes, 117, 124, 127; conchatus, 126,  
127; densus, 126, 127; fulvus, 126,  
127; ignarius laevigatus, 119, 126;  
ignarius resupinatus, 120; nigro-  
limitatus, 126; Pini, 126, 127; pute-  
arius, 126, 127; robustus, 122; sali-  
cinus, 120
- Fomitiporia laminata, 119; obliqui-  
formis, 119; prunicola, 118; tsugina,  
121; Weirii, 122
- Fragaria, 304
- Fraxinus, 120
- Further study of the morphology and  
life history of the rose black spot  
fungus, B. O. Dodge, 446
- Fusarium, 53, 291, 302; Batatatis,  
304; conglutinans, 302; congluti-  
nans Callistephi, 301; hyperoxyspo-  
rum, 304; lagenarium, 138; Lyco-  
persici, 304; Martii Phaseoli, 302;  
niveum, 304; oxysporum, 304; va-  
sinfectedum, 302, 303
- Fuscoporia, 127
- Fusisporium Rubi, 302

- Geotrichum candidum*, 314  
*Geranium*, 358, 359; *Berberianum*, 357; *chilloense*, 359; *Core-Core*, 358; *Ochsenii*, 358; *Sodiironum*, 359  
*Gloeodes pomigena*, 301  
*Gloeosporium*, 134, 159-161, 167-169, 171, 175, 180; *Camphorae*, 302; *fructigenum*, 171, 175, 177; *lagenarium*, 138; *reticulatum*, 138; *Rosae*, 223, 304; *rufomaculans*, 165, 168, 171, 175, 177  
*Gloeosporium Rosae*, a nomen nudum, Anna E. Jenkins, 223  
*Glomerella*, 161, 182, 223; *cincta*, 223; *cingulata*, 136, 138, 301, 303; *Gossypii*, 302; *lagenaria*, 138; *rufomaculans*, 168, 175-177, 182, 185  
*Glycyrrhiza lepidota*, 82  
*Gnomonia Caryae*, 303; *ulmea*, 302  
*Godetia*, 485; *Cavanillesii*, 485  
*Gossypium*, 478; *hirsutum*, 302  
*Gouania*, 475; *Blanchetiana*, 475; *corylifolia*, 474; *polygama*, 474; *pyrifolia*, 475  
*Graphiola Phoenixis*, 303  
*Grindelia texana*, 79  
*Griselinia ruscifolia*, 490  
*Guignardia Bidwellii*, 303  
*Gymnoconia*, 97; *interstitialis*, 302  
*Gymnolomia multiflora*, 79  
*Gymnosporangium bermudianum*, 302; *Betheli*, 78; *globosum*, 301; *Juniperi-virginianae*, 301  
*Gyromitra esculenta*, 409  
  
*Hainesia*, 458  
*Hamaspora*, 338, 340, 341  
*Haplopyxis Crotalariae*, 346  
*Haploravenelia Ingae*, 336  
*Helianthus fascicularis*, 79; *lenticularis*, 79  
*Helminthosporium*, 390; *Sacchari*, 304  
*Helotium Seaveri*, 247-249  
*Hemibradyspora*, 260, 262, 268, 270, 292  
Herbarium arrangement of mycological specimens, E. T. Bartholomew, 227  
*Heteropteris*, 361, 364  
*Heterosporium gracile*, 303  
*Hexagonia*, 126  
*Hibiscus esculentus*, 303, 379, 394; *syriacus*, 303; *tiliaceus*, 379, 382  
*Hicoria*, 303; *Pecan*, 303  
*Hieracium scabriusculum*, 80  
*Holcus Sorghum*, 304  
*Holwayella*, 104  
*Hordeum jubatum*, 81  
*Horkelia*, 435  
*Hura crepitans*, 373  
  
*Hydrocotyle*, 486, 487; *aconitifolia*, 486; *bonariensis*, 486; *Bonplandii*, 487; *Poeppigii*, 486; *umbellata*, 487; *Volkmanni*, 487  
*Hymenocallis*, 380; *crassifolia*, 380; *littoralis*, 380  
*Hypera punctata*, 412  
*Hypericum*, 481; *brasiliense*, 481; *chilense*, 481  
  
*Icacorea*, 490  
*Ilex*, 125, 303  
*Indigofera arrecta*, 290; *suffruticosa*, 350  
*Inga*, 333, 336; *edulis*, 336; *insignis*, 336  
Inheritance of the albinistic non-conidial character in interspecific hybrids in *Neurospora*, B. O. Dodge, 1  
*Ipomoea*, 495, 496, 498, 499; *acuminata*, 495; *Batatas*, 267, 273, 275-277, 279, 281, 283, 304; *biloba*, 399; *fistulosa*, 497; *Pes-tigridis*, 399; *polymorpha*, 496; *purpurea*, 399, 495  
*Iresine calea*, 404; *paniculata*, 404  
*Iris*, 303  
*Irpex*, 130; *deformis*, 130; *fimbriiformis*, 132; *griseofuscus*, 131; *hirsutus*, 131, 132; *lacteus*, 130, 131; *obliquus*, 130, 132; *paradoxus*, 130  
*Iva axillaris*, 80  
*Ivesia*, 435, 436; *Gordonii*, 435  
  
Jackson, H. S., The rusts of South America based on the Holway Collections—III, 96; IV, 332; V, 463  
*Jacquemontia*, 499; *ferruginea*, 498; *punctantha*, 499  
Jenkins, Anna E., *Gloeosporium Rosae*, a nomen nudum, 223  
*Juncus Dudleyi*, 82  
*Juniperus*, 248; *communis*, 248, 250; *virginiana*, 302  
  
*Kalmia latifolia*, 303  
Kelley, Arthur Pierson, The concept of *Mycorhiza*, 147  
Kern, F. D., and Thurston, H. W., Jr., Notes on some rust collections from Colorado, Wyoming, and South Dakota, 77  
*Kuehneola*, 105, 106; *andicola*, 105; *Arthuri*, 106; *Gossypii*, 478; *Loeseneriana*, 105, 106; *malvicola*, 303, 478  
  
*Lachnea*, 313  
*Lactuca pulchella*, 79; *sativa*, 303  
*Lafoensia Pacari*, 483  
*Lagerstroemia australiana*, 302  
*Lahaina*, 273, 275, 277, 279, 281, 283

- Lantana, 375  
 Lathraea, 147  
 Lathyrus, 82; magellanicus, 354; odoratus, 304  
 Lavauxia, 485; mutica, 485  
 Lenzites, 130  
 Leonurus Cardiaca, 396  
 Leptomitius, 308  
 Leptosphaeria Coniothyrium, 302, 304  
 Leptothyrium carpophilum, 303  
 Leucaena microphylla, 337  
 Ligusticum Porteri, 80; simulans, 80  
 Ligustrum, 304  
 Lisianthus elegans pedunculatus, 491  
 List of diseases of economic plants in Alabama, W. L. Blain, 300  
 Lonchocarpus, 350  
 Luehea, 477  
 Lupinus, 345, 352; argenteus, 82; paniculatus, 345, 352; soratensis, 352  
 Lycopersicon, 184; esculentum, 180, 304, 390  
 Maclura, 120  
 Macrosporium cucumerinum, 302; herculeum, 304; Nerii, 303; Tomato, 304  
 Magnolia, 303  
 Mahonia, 359  
**Mainsia**, 106; *chinensis*, 107; *clara*, 108, 112, 113; *cundinamarcensis*, 108, 114; *epiphylla*, 108, 112; *Holwayii*, 107, 109, 116; *Lagerheimii*, 108, 110; *Mayorii*, 108, 112, 115; *peruviana*, 107, 108; *Pittieriana*, 107, 110; *quitensis*, 107, 108, 113, 115, 116; *Rubi*, 108, 110, 112; *Rubi-urtici*, 107; *Rubi-urticifolia*, 108; *tenella*, 108, 111; *urediniformis*, 107; *urticifolii*, 115; *variabilis*, 108, 111  
 Malus sylvestris, 301  
 Malva, 480; parviflora, 480; pusilla, 81  
 Malvastrum, 479, 480; capitatum, 478; coromandelianum, 480  
 Malvaviscus, 478  
 Mandevilla Mandoni, 492  
 Manihot, 382, 472; utilissima, 371, 382, 404  
 Marssonina, 456; Martini, 303  
 Medicago Medicaginis, 302; sativa, 301, 354  
 Meibomia, 354; Scorpiurus, 351  
 Melampsora argentinensis, 465; Bigelowii, 78; Humboldtiana, 78  
 Melampsoropsis Pyrolae, 78  
 Mentha Penardi, 81  
 Mentzelia, 482  
 Mesosetum ferrugineum, 297; lolii-forme, 298  
 Microascus, 314-317, 321-325, 327-329; intermedius, 327; longirostris, 322; sordidus, 315, 322, 328; **trigonosporus**, 317, 319, 322, 323, 326-328; *variabilis*, 315, 323  
 Micropuccinia Arechavelatae, 473; heterospora, 479; Leveillei, 358  
 Microsphaera Alni, 303  
 Microsporium, 88; fulvum, 88  
 Mikania, 398  
**Mimema**, 338, 340, 341; **Holwayi**, 338, 340, 348, 349  
 Mimosa albida, 337; sepiaria, 335  
 Mitella, 80  
 Modiola multifida, 388  
 Monarda Ramaleyi, 81  
 Monilia, 47, 48, 313; candida, 317; sitophila, 48  
 Mortierella, 308  
 Morus, 303  
 Mucor, 33, 89; parasiticus, 307  
 Mucronoporus fulvidus, 117, 118  
 Mulinum spinosum, 487  
 Musa sapientum, 273, 276, 277  
 Mycorhiza, 147  
 Mycosphaerella Fragariae, 304  
 Myrcia, 484  
 Myriangium, 303; tuberculans, 303  
 Nasella, 479  
 Nectandra oppositifolia, 103  
 Nematosporangium, 252-260, 262-265, 267, 268, 290, 291, 294; aphanidermatum, 259, 262, 264, 269, 271, 284-286; aphanidermatum **hawaiiensis**, 286, 287; **arrhenomanes**, 264, 269, 270, 272, 273, 292; **arrhenomanes hawaiiensis**, 269, 270; **Butleri**, 259, 262, 264, 269, 271, 286, 288; dictyosporum, 191, 199; **epiphanosporon**, 264, 269, 271, 282, 283, 292, 293; **hyphalosticton**, 269, 270, 275, 293; **Indigoferae**, 262, 269, 271, 289, 290, 294; **leiohyphon**, 269, 271, 282, 283, 293; **leucosticton**, 269, 271, 281, 282, 293; **polyandron**, 269, 270, 276, 278, 293; **rhizophthoron**, 269, 270, 271, 279, 280, 293; **spaniogamon**, 269, 270, 273, 276; **thysanohyphalon**, 269, 270, 277  
 Nemopanthes, 125  
 Neanelsonia acuminata, 487  
 Nephrospora, 317, 329; Mangini, 317  
 Nerium Oleander, 303  
 Neurospora, 8, 18, 44, 47, 313; crassa, 1-4, 7, 8, 31, 45; sitophila, 1-5, 7-14, 22, 26, 28, 30, 31, 33, 34, 45, 46, 48; tetrasperma, 1, 2, 9-13, 15-17, 19, 20, 24, 27-37, 41-43, 45-47, 49, 50  
 Nicotiana, 368; repanda, 387; Tabacum, 368, 387



- Nigredo Fabae, 353; Medicaginis, 354; Neurocarpi, 355  
 Notes on Iowa species of the genus *Irpex*, K. Cejp, 130  
 Notes on new species of Ustilaginales, George L. Zundel, 296  
 Notes on some rust collections from Colorado, Wyoming, and South Dakota, H. W. Thurston, Jr., and F. D. Kern, 77  
 Nyssopsora, 339  
  
 Observations on *Achorion gypseum*, C. W. Emmons, 87  
 Observations on *Pythium dictyosporum*, Frederick J. Sparrow, Jr., 191  
 Occurrence of *Schizosphyllum commune* on green apples, F. D. Bailey and S. M. Zeller, 154  
*Odosstemon aquifolium*, 81  
*Oedogonium*, 59  
*Oenothera*, 485; *mollissima*, 485  
*Oligandra*, 260, 262, 264, 271, 291, 292-294  
*Oligocomba*, 260, 262, 271, 293  
*Olivia capituliformis*, 466  
*Operculina glabra*, 495  
*Opuntia sulphurea*, 482  
*Oreobatus*, 436; *deliciosus*, 78  
*Orobancha*, 147  
*Osmorrhiza*, 489; *Berteri*, 489  
*Ostrya*, 120  
 Overholts, L. O., Diagnoses of American *Porias*—III. Some additional brown species, with a key to the common brown species of the United States and Canada, 117  
*Oxalis*, 359; *scandens*, 359  
*Oxydendrum*, 120  
*Oxypolis Fendleri*, 80, 81  
*Oxystelma*, 494  
  
*Paeonia*, 303  
*Panicum barbinodum*, 273, 275, 277, 279, 281, 283, 284; *demissum*, 298; *missionum*, 298; *revulare*, 296  
*Papulaspora*, 310  
*Parasitella*, 307, 308; *simplex*, 307  
*Parosela pagensis*, 357  
*Paspalum millegranum*, 299  
*Passalora penicillata*, 370  
*Passiflora*, 381; *incarnata*, 381; *sexiflora*, 381  
*Paulownia tomentosa*, 304  
*Pavonia rosea*, 479; *sepium*, 478; *speciosa*, 478; *spinifex*, 478  
*Pelargonium*, 303  
*Penicillium*, 160, 313-315, 322, 323, 326, 329; *brevicaule*, 315  
*Pennisetum*, 299  
*Pereskia*, 472; *grandifolia*, 472, 473  
  
*Peridermium*, 233, 240; *Ipomoeae*, 495  
*Peristomium*, 317, 325, 326, 329; *desmosporum*, 316, 317, 325; *desmosporum Verticillium*, 325  
*Pernettya Pentlandii*, 490  
*Peronoplasmopara cubensis*, 302, 304  
*Peronospora*, 201; *parasitica*, 302  
*Persea americana*, 301; *gratissima*, 266, 372; *palustris*, 372  
*Pestalozzia Guepini*, 303  
*Petunia parviflora*, 390  
*Peziza chloromela*, 247, 250  
*Pezizella Lythri*, 458  
*Phakopsora*, 347, 465; *argentinensis*, 465; *Crotalariae*, 346; *Psoraleae*, 346; *Vitis*, 476  
*Phaseolus*, 351, 384, 390; *aureus*, 273, 275, 276, 277; *coccineus*, 302; *lunatus*, 384, 390; *Mungo*, 384; *Soja*, 267; *vestitus*, 351; *vulgaris*, 302, 384, 390  
*Philibertia*, 494  
*Phleum pratense*, 81  
*Phlox*, 303  
*Phoenix*, 303  
*Phoma*, 453, 456; *carpogena*, 456; *destructiva*, 303; *Lingam*, 302  
*Phomopsis Citri*, 302; *juniperovora*, 302; *vexans*, 302  
 Photographs and descriptions of cup-fungi—XIV. A new genus, Fred J. Seaver, 247; XV. The giant *Elvela*, 409  
*Phragmidium*, 97, 433, 434, 437, 441, 444; *alaskanum*, 436, 442, 443; *americanum*, 438, 440-443; *Andersoni*, 436, 441, 442; *biloculare*, 435, 442, 443; *disciflorum*, 104, 438-440, 442, 443; *Horkeliae*, 435, 441, 443; *Ivesiae*, 78, 435, 442, 443; *Jonesii*, 436, 441, 443; *montivagum*, 78, 438-443; *occidentale*, 437, 442, 443; *Peckianum*, 78, 436, 437, 442, 443; *Potentillae*, 434, 435, 441, 443; *Rosae-acicularis*, 438-443; *Rosae-arkansanae*, 438-443; *Rosae-californicae*, 439, 440, 442, 443; *Rosae-pimpinellifoliae*, 438-440, 442, 443; *Rosae-setigerae*, 438; *Rubi-idaei*, 434, 435, 437, 442, 443; *Rubiodorati*, 435, 437, 442, 443; *speciosum*, 78, 437, 438, 442, 443; *subcorticium*, 439  
*Phragmidium* species of North America: differential teliospore and aecial characters, George B. Cummins, 433  
*Phycomyces Pirotianus*, 308  
*Phyllactinia corylea*, 302  
*Phyllanthus brasiliensis*, 466, 468; *lathyroides*, 469



- Phyllosticta acericola, 303; Batatas, 304; camelliicola, 303; Caryae, 303; Cookei, 303; kalmicola, 303; micro-punctata, 301; Mucunae, 304; Paulowniae, 304; pyrina, 303; solitaria, 301  
 Phyllostictina, 456; carpogena, 453  
 Physalospora Cydoniae, 301; Zeae, 245  
 Physoderma Zeae-maydis, 302  
 Physopella Fici, 303; Vitis, 476  
 Phytolacca decandra, 400; icosandra, 400  
 Phytanomeris punctatus, 412  
 Phytophthora, 253-257, 291, 308, 310; cactorum, 303; infestans, 304  
 Picea, 121  
 Pieris, 490; Brassicae, 411, 412  
 Pilobolus, 308  
 Pinus, 121; echinata, 304; palustris, 303; Taeda, 303  
 Piptadenia, 333, 334; latifolia, 332, 333; laxa, 332  
 Pipturus albidus, 383; Gaudichaudianus, 383  
 Plantago, 392; lanceolata, 392; lusitanica, 392; major, 392  
 Plasmopara, 255  
 Plectodiscella veneta, 302  
 Pleoravenelia Indigoferae, 350  
 Plethorocomba, 260, 262, 271, 293  
 Poreia psoraloides, 348  
 Polyandra, 260, 262, 264, 268, 269, 291, 292, 294  
 Polyporus, 117, 124, 240; gilvus, 126; glomeratus, 125, 126, 128  
 Polystigma, 44  
 Polythelis Thalictri, 78  
 Pomolobus pseudoharengus, 412  
 Poria, 117, 118, 123, 124, 127, 128; betulina, 126; contigua, 120; ferrea, 117, 126-129; ferrugineo-fusca, 126, 127; ferruginosa, 120, 126, 127; Friesiana, 119, 120; fulvida, 127; inermis, 118, 125; laminata, 119, 120, 127; Macounii, 127; marginella, 127; mucida, 130; obliqua, 120; obliquiformis, 121, 127; prunicola, 118, 126-128; punctata, 119, 125, 127-129; subiculosa, 125, 128; superficialis, 128; tsugina, 121, 125, 128, 129; viticola, 128; Weirii, 122, 126, 128, 129  
 Porophyllum ruderales, 394  
 Potentilla, 435; glaucophylla, 78  
 Prunus, 119, 126, 127; americana, 304; domestica, 304; Persica, 105  
 Pseudocolus, 84; Schellenbergiae, 84  
 Pseudomonas campestris, 302; Malvacearum, 302  
 Psidium, 484; Guajava, 484  
 Psoralea argophylla, 82; glandulosa, 347; lasiostachya, 347  
 Ptilidium, 458  
 Puccinia, 97, 98, 100, 104, 348, 495; Absinthii, 79; Actinostemonis, 466; aemulans, 79; Amiciae, 356; Antirrhini, 304; Araujae, 493; Archavelatae, 473; Arenariae, 304; Asterum, 79; Banisteriae, 362; Berberidis, 96; Berberidis-Darwinii, 96; Bergii, 348; Berkeleyana, 496; Boissduvaliae, 485; calliquensis, 357; Cirsii, 79; Clematidis, 79; Cleomis, 103; compacta, 493; conrescens, 493; Convolvuli, 495; coronata, 303; crassipes, 495; cruciferarum, 79; Cynanchi, 494; Daleae, 357; Dichondrae, 496, 497; discors, 487; distenta, 357; distinguenda, 497; enecta, 498; Epilobii-tetragoni, 485, exilis, 479; festata, 467; Gayophyti, 485; Gentianae, 79; Geranii, 358; Geranii-silvatici, 358; geraniicola, 358; Gonolobi, 494; Gouaniae, 474; Grindeliae, 79; Griseliniae, 490; Grossulariae, 79; grumosa, 79; Hedysari-paniculati, 353; Helianthi-mollis, 79; hemisphaerica, 79; heterogena, 479; Heteropteridis, 361; heterospora, 479; invaginata, 474; Ipomoeae, 496; Ipomoeae-panduratae, 496; Jambosae, 483; Jambulana, 484; Kunzeana, 494; Leveilleana, 358; Leveillei, 358; Ligustici, 80; Lisianthi, 491; Lithospermi, 497; luxurians, 485; Malvacearum, 81, 480; Mandevillae, 492; Menthae, 81; montanensis, 101; mundula, 488; munita, 496; neurophila, 484; nocticolor, 497; obliqua, 492, 494; obscurata, 487; offuscata, 350; opulenta, 498; Opuntiae, 482; oxalidis, 359; Oxypetalis, 494; papillifera, 332; paraensis, 475; Peckii, 81; Philibertiae, 494; Philippii, 489; picturata, 363; Piptadeniae, 333, 348; platyspora, 480; poculiformis, 81; Polygoni-vivipari, 81; Pruni-spinosae, 105; Psidii, 483, 485; purpurea, 304; repentina, 489; scaber, 81; scandica, 81; Schnyderi, 493; Senecionis, 97; Sherardiana, 81, 480; Sorghi, 451; Sphaerostigmatis, 485; subneurophila, 484; Tabernaemontanae, 492; Tournefortiae, 501; tournefortiicola, 501; Treleasiana, 81; Trifolii, 355; triti-

- cina, 304; tuyatensis, 497, 498; urticata, 81; Viola, 81, 482; Xanthii, 81; Zorniae, 350
- Pucciniastrum Agrimoniae, 79, 104; Myrtilli, 490
- Puccinosira, 476, 477; pallidula, 476, 477; Triumphetae, 477
- Pyrethrum, 204
- Pyrola minor, 78
- Pyronema, 8, 33, 44
- Pyrus communis, 303
- Pythiogeton, 309
- Pythiosis, 309, 310
- Pythium, 191, 196, 197, 199, 201, 253-258, 260, 290, 291, 308, 310; aphanidermatum, 254, 284; arrhenomanes, 201, 272; Butleri, 286; complens, 254; dictyosporum, 191, 194, 196, 197, 199, 200-202; gracile, 254; Indigoferae, 290; monospermum, 254; reptans, 254
- Quercus, 118, 120, 131, 132; macrocarpa, 131; nigra, 303
- Rafflesia, 147
- Ragnhildiana**, 365, 402; **Agerati**, 402, 403; **Cyathulae**, 403; **gonatoclada**, 403; **Manihotis**, 404; **Tremae**, 405
- Ranunculus alismellus, 82
- Rapanea, 491; pseudocrenata, 491; umbellata, 491
- Rare phalloid from The New York Botanical Garden, Fred J. Seaver, 83
- Ravenelia, 339, 466; Acaciae-Farnesianae, 335; appendiculata, 466, 468; capituliformis, 466; echinata, 334; ectypa, 334; **faceta**, 342; Henningsiana, 334; Hieronymi, 335; **idonea**, 335; Indigoferae, 350; Ingae, 335; **irregularis**, 336; Lagerheimiana, 337; Leucaenae-microphyllae, 337; Lonchocarpi, 350; microcarpa, 343; microspora, 343; Mimosae, 335; Mimosae-albidae, 337; platensis, 345; **rata**, 337
- Rheosporangium aphanidermatum, 284, 310
- Rhizoclonium hieroglyphicum, 193, 200
- Rhizoctonia Solani, 304
- Rhododendron, 302
- Rhopobota, 421; vacciniiana, 412, 425
- Rhytisma ilicincola, 303
- Ribes albiflorum, 103; americanum, 79; inebrians, 78
- Ricinus communis, 390
- Rivina humilis, 400
- Robinia, 120
- Roestelia interveniens, 478
- Rosa, 78, 104, 223, 304; acicularis, 438; alba, 438; arkansana, 438; californica, 439; carolina, 438; eglanteria, 438; Engelmanni, 438; Fendleri, 438, 441; gallica, 438; gymnocarpa, 439; hemisphaerica, 438; nutkana, 438, 439, 441; setigera, 439; suffulta, 78, 438; virginiana, 438
- Rostrupia praelonga, 478
- Rubus, 105, 106, 109, 110, 112-114, 223, 302, 304, 341; bogotensis, 110, 112, 115; boliviensis, 114; brasiliensis, 106; erythroclados, 106; floribundus, 106, 109, 110, 115; glaucus, 110; megalococcus, 111; nubigenus, 111; odoratus, 437; parviflorus, 437; peruvianus, 114; roseus, 113; stellatus, 436; strigosus, 437; trivialis, 112; urticaefolius, 106, 109, 110; urticifolius, 115
- Rusts of South America based on the Holway Collections—III, H. S. Jackson, 96; IV, 332; V, 463
- Sabouraudites, 88
- Saccharomyces Freseniusi, 140; roseus, 140
- Saccharomycodes Ludwigii, 74
- Saccharum officinarum, 273, 275, 277, 279, 281, 283, 284, 304
- Salix, 78, 120; amygdaloides, 78
- Sambucus, 370; canadensis, 370; mexicana, 370; nigra, 370; nigra laciniata, 370; racemosa, 370
- Sapium, 471
- Saprolegnia, 309
- Sawyer, Wm. H., Jr., Studies on the morphology and development of an insect-destroying fungus, Entomophthora sphaerosperma, 411
- Schizoparme straminea, 457
- Schizophyllum commune, 154
- Schizosaccharomyces octosporus, 33
- Schroeteria aster argentinensis, 465; Ulei, 466, 469
- Sclerospora, 255, 310
- Sclerotinia, 313; carunculoides, 303; fructicola, 303, 304; sclerotiorum, 303
- Sclerotium, 458
- Sclerotium, 219; bataticola, 304; Delphinii, 204, 206, 208-219, 221, 222; Rolfsii, 204-208, 210, 212-216, 218-222, 302-304
- Scopulariopsis, 314-317, 319, 323, 325, 328, 329; candida, 317; cinerea, 314, 316, 317; **trigonospora**, 317, 328
- Seaver, Fred J., A rare phalloid from The New York Botanical Garden, 83; Photographs and descriptions

- of cup-fungi—XIV. A new genus, 247; XV, The giant Elvela, 409
- Secale cereale*, 304
- Septobasidium*, 301–303; pedicellatum, 301, 302; retiforme, 303
- Septoria divaricata*, 303; *Lycopersici*, 304; *nodorum*, 304
- Sequoia*, 127, 247, 248; *sempervirens*, 249, 250
- Serjania*, 474; *cuspidata*, 473
- Sida cordifolia*, 479; *paniculata*, 480; *rhombofolia*, 481; *spinosa*, 480; *tomentosa*, 480; *urens*, 480
- Sideris*, C. P., Taxonomic studies in the family Pythiaceae—I. *Nematosporangium*, 252
- Soja* Max, 304
- Solanum* *Melongena*, 302; *nigrum*, 384; *tuberosum*, 184, 267, 273, 275, 276, 277, 279, 281, 283, 284, 304
- Solheim, W. G., and Stevens, F. L., *Cercospora* studies—II. Some tropical *Cercosporae*, 365
- Sordaria*, 323
- Sparrow, Frederick K., Jr., Observations on *Pythium dictyosporum*, 191
- Spermocoe ocymoides*, 385
- Sphacelotheca braziliensis*, 297; **Chasseae**, 297; *cruenta*, 304; **echinata**, 298; *Holwayi*, 297; *Kellermanii*, 297; *leucostachys*, 297; **Mesoseti**, 298; **Vryburgii**, 298
- Sphaeralcea*, 479, 480; *coccinea*, 81; *obtusifolia*, 479
- Sphaerocarpos*, 47; *Donnellii*, 2
- Sphaerosporangium*, 253–255, 308
- Sphaerostigma*, 486; *tenuifolium*, 485
- Sphaerotherca*, 8, 44, 235; *pannosa*, 304
- Sphenospora*, 98; *Berberidis*, 99; *quintensis*, 99
- Spirechina*, 105–107; *Arthuri*, 105, 106; *chinensis*, 107; *cundinamaricensis*, 114; *epiphylla*, 112; *Loeseneriana*, 105; *Pittieriana*, 110, 114; *Rubi*, 110, 114; *variabilis*, 111
- Spirogyra*, 191–193, 196, 200; *crassa*, 191, 200, 202; *insignis*, 191; *nitida*, 191, 200
- Spongospora subterranea*, 304
- Stemphylium*, 182, 183
- Stevens, F. L., A comparative study of *Sclerotium Rolfsii* and *Sclerotium Delphinii*, 204; The ascigerous stage of *Colletotrichum lagenarium* induced by ultra-violet irradiation, 134
- Stevens, F. L., and Solheim, W. G., *Cercospora* studies—II. Some tropical *Cercosporae*, 365
- Stigmaphyllon*, 361, 363; *acuminatum*, 362; *affine*, 362; *tomentosum*, 362; *vitifolium*, 362
- Stipa*, 479; *Lettermanii*, 81
- Stizolobium*, 304
- Studies on *Ascoidea rubescens*—I. History and development, Leva B. Walker, 51
- Studies on the morphology and development of an insect-destroying fungus, *Entomophthora sphaerosperma*, Wm. H. Sawyer, Jr., 411
- Symphoricarpos albus laevigatus*, 159, 178; *orbiculatus*, 164
- Syringa vulgaris*, 303
- Tabernaemontana amygdalifolia*, 492
- Tachyspora*, 260, 262, 263, 270, 292, 293
- Taphrina coerulescens*, 303; *deformans*, 303; *epiphylla*, 33; *Robinsoniana*, 301
- Taraxacum*, 80; *Taraxacum*, 80
- Tarichium*, 411; *sphaerospermum*, 412
- Taxonomic studies in the family Pythiaceae—I. *Nematosporangium*, C. P. *Sideris*, 252
- Tectona grandis*, 399
- Thalictrum*, 79; *dasycarpum*, 78
- Themeda Forskalii*, 298
- Thermoascus aurantiacus*, 316
- Thielavia*, 313, 314, 323, 329; *basicola*, 313; *terricola*, 313
- Thraustotheca*, 196
- Thuja*, 248, 250; *occidentalis*, 248, 249; *plicata*, 123, 249
- Thurston, H. W., Jr., and Kern, F. D., Notes on some rust collections from Colorado, Wyoming, and South Dakota, 77
- Tilletia Paspali*, 299; **transvaalensis**, 299
- Tolypothrix*, 193, 200
- Torula*, 141; *corallina*, 144; *glutinis*, 140; *minuta*, 145; *rufa*, 145
- Torulopsis*, 144, 145; *Biourgei*, 144; *corallina*, 144; *minuta*, 145, 146; *minuta parvissima*, 146; *Montii*, 144; *rufula*, 145; *saccharophoba*, 144; *Saitoi*, 144
- Tournefortia*, 501–503; *brachiata*, 500; *fuliginosa*, 501; *grandifolia*, 501; *psilostachya*, 502; *sericea*, 502; *suaveolens*, 502
- Trametes*, 117, 125; *carbonaria*, 125, 126, 128; *odorata*, 125; *Sequoiae*, 127, 128; *setosus*, 128; *tenuis*, 120, 125, 126, 128
- Tranzschelia punctata*, 104
- Trema micrantha*, 405
- Triactella*, 339, 341, 342; **Holwayi**, 341, 342; *pulchra*, 342

- Trichobasis euphorbiaecola, 471;  
 Howe, 494; Hydrocotyles, 486  
 Trichophyton asteroides, 94; gyp-  
 seum, 92  
 Trichopsora Tournefortiae, 503  
 Trifolium, 82, 355, 356; amabile, 355;  
 Parryi, 82; pratense, 302; repens,  
 82, 353, 356  
 Triphragmiopsis, 339  
 Triphragmium, 339, 341, 342  
 Triticum aestivum, 304; vulgare, 273,  
 275, 279, 281, 283, 284  
 Triumfetta, 477; longicornis, 476;  
 semitriloba, 477  
 Tsuga, 121, 122  
  
 Ulmus, 302  
 Uncinula, 302; australiana, 302  
 Uredo, 104, 347, 360, 465, 468, 469,  
 483; Andromedae, 490; Anilis, 350;  
 Apocynaceae, 493; appendiculata,  
 351; asclepiadina, 494; Asclepiadis,  
 494; banistericola, 362; Betae Con-  
 volvuli, 495; bonariensis, 486; capi-  
 tuliformis, 466; cassicola, 342;  
 Cisnerosana, 470; **Condylocarpi**, 493;  
 Cornui, 467; Crotalariae, 346; cro-  
 tonicola, 465; Crotonis, 465, 468,  
 470; Cupheae, 483; **cupheicola**, 482;  
 Cynapii, 487; Desmodii-leiocarpi,  
 353; Dieteliana, 498; **emendata**,  
 350; **Eriosemae**, 351; Erythroxylo-  
 nis, 360; Eugeniurum, 483; excipu-  
 lata, 336; Fabae, 353; flavidula,  
 483; floridana, 482; Goeldiana, 483;  
 Gossypii, 478; Gouaniae, 474; Hy-  
 drocotyles, 486; imperialis, 105; In-  
 gae, 335; Ipomoeae, 495; Ipomoeae-  
 pentaphyllae, 495; **Lafosseae**, 483;  
 Leuhae, 477; Loeseneriana, 105;  
 malvicola, 478; medicaginicola, 354;  
 mulinicola, 487; Myrciae, 483; Myr-  
 tacearum, 483; neurophila, 483;  
 notata, 361; ochraceo-flava, 110;  
 Oxalidis, 359; pachystegia, 502;  
 Pamparum, 351; **pavida**, 468; Phyl-  
 lanthi, 466, 469; proëminens, 471;  
 rhombica, 472; rufa, 351; **seclusa**,  
 484; solitaria, 351; spinulosa, 498,  
 499; subneurophila, 483; Tijuae,  
 469; **Tournefortiae**, 502; unilater-  
 alis, 359; vagans Epilobii-tetragoni,  
 485; **valentula**, 469; Vitis, 476;  
**Zarumae**, 481; Zorniae, 350  
 Uromyces, 103, 105, 345, 463, 473;  
**Actinostemonis**, 470; aemulus, 82;  
 aeruginosus, 473; andinus, 110, 463,  
 470; appendiculatus, 302, 351, 356;  
 Argophyllae, 82; Arthuri, 106; As-  
 clepiadis, 494; **Bradburyae**, 352;  
 brasiliensis, 498, 499; castaneus,  
 352; Chilensis, 354; Cisnerosanus,  
 470; clavatus, 354; cundinamarcen-  
 sis, 114; Desmodii-leiocarpi, 353;  
 Dieteliana, 343; dolichosporus,  
 502; elatus, 352; elegans, 356; Eu-  
 phorbiae, 471; euphorbicola, 471;  
 Fabae, 82, 353; flectens, 353; foveo-  
 latus, 343; gemmatus, 498; gigan-  
 teus, 498, 499; Glycyrrhizae, 82;  
 Hedysari-obscuri, 97; Hedysari-  
 paniculati, 345, 353; Hemmen-  
 dorffii, 343; heterodermus, 82;  
 Howe, 494; Hyperici-frondosi, 481;  
 hypsophilus, 463, 470; ingicola, 336;  
 insularis, 354; Jonesii, 82; Lager-  
 heimii, 110; lathyrinus, 354; Lo-  
 esenerianus, 105; malvacearum, 479;  
 malvicola, 479; Medicaginis, 354;  
 Mulini, 487; Mulini magellanica,  
 487; Myrsines, 490; nerviphila, 353;  
 Neurocarpi, 354; oblongus, 82, 356;  
 orbicularis, 355; pachycephalus,  
 481; **Pereskiae**, 473; Perlebiae, 343;  
 pervius, 473; Pittierianus, 110;  
 platysporus, 480; porcensis, 336;  
 proëminens, 471; quitensis, 112,  
 115; Rhapanea, 490, **ribicola**, 103;  
 rostratus, 354, 355; Rubi, 110;  
 Rubi-urticifolii, 115; Scirpi, 486;  
 Silphii, 82; substriatus, 82; tenui-  
 stipes, 355; **tolerandus**, 471; Tour-  
 nefortiae, 502; Trifolii, 82, 353, 355;  
**Trifolii-megalanthi**, 356; Trifolii-  
 repentis, 353; urediniformis, 107;  
 Usterianus, 490, 491; Usterii, 105;  
 variabilis, 111; **verus**, 344; **vicinus**,  
 499; Vignae, 302, 356  
 Urophlyctis Alfalfae, 301  
 Uropyxis **Amiciae**, 356; Amorphae,  
 81; Crotalariae, 346; Daleae, 357;  
 quitensis, 97, 99; sanguinea, 81  
 Ustilago, 297; Avenae, 303; **brazili-  
 ensis**, 296; **gregaria**, 296; olivacea  
 Pseudocyperi, 297; Tritici, 304;  
 Zeae, 302  
  
 Vaccinium, 78; macrocarpon, 412  
 Valsa nivea, 245; sordida, 245  
 Venenarius Lanei, 226  
 Venturia inaequalis, 301; pyrina, 303  
 Verbena, 204  
 Vermicularia depazeoides, 370  
 Verticillium, 291, 303  
 Vicia Faba, 275, 353  
 Vigna Catjang, 384, 390; glabra, 390;  
 luteola, 356; sinensis, 302  
 Viola, 482; adunca, 81; maculata pu-  
 bescent, 482; odorata, 304  
 Vitis, 303, 476  
 Volvaria, 152, 153; gloiocephala, 153;  
 speciosa, 152, 153  
 Volvaria speciosa, John Dearness, 152

- Walker, Leva B., Studies on *Ascoidea rubescens*—I. History and development, 51  
*Willia anomala*, 59; *Saturnus*, 74  
*Wissadula hernandioides*, 480
- Xanthium speciosum*, 81  
*Xanthoxylum*, 364  
*Xolisma*, 490  
*Xylaria*, 301
- Zea Mays*, 265, 273, 275, 277, 281, 283, 302  
Zeller, S. M., *Amanita calyptrata* and *Amanita calypetroderma*, 225  
Zeller, S. M., and Bailey, F. D., The occurrence of *Schizophyllum commune* on green apples, 154  
Zinnia, 389  
*Zornia diphylla*, 350  
Zundel, George L., Notes on new species of *Ustilaginales*, 296